



Transverse (Spin) Structure of Hadrons


Matthias Burkardt

burkardt@nmsu.edu

New Mexico State University


Outline

- Probabilistic interpretation of GPDs as Fourier trafos of impact parameter dependent PDFs
 - $H(x, 0, -\Delta_{\perp}^2) \longrightarrow q(x, \mathbf{b}_{\perp})$
 - $\tilde{H}(x, 0, -\Delta_{\perp}^2) \longrightarrow \Delta q(x, \mathbf{b}_{\perp})$
 - $E(x, 0, -\Delta_{\perp}^2) \longrightarrow \perp$ distortion of PDFs when the target is \perp polarized



- Chromodynamik lensing and \perp SSAs

transverse distortion of PDFs
+ final state interactions } $\Rightarrow \perp$ SSA in $\gamma N \longrightarrow \pi + X$




- $g_2(x) \leftrightarrow$ transverse force on quarks in DIS
- transversity distribution in an unpolarized target
- Summary

Generalized Parton Distributions (GPDs)

- GPDs: decomposition of form factors at a given value of t , w.r.t. the average momentum fraction $x = \frac{1}{2} (x_i + x_f)$ of the active quark

$$\int dx H_q(x, \xi, t) = F_1^q(t) \quad \int dx \tilde{H}_q(x, \xi, t) = G_A^q(t)$$
$$\int dx E_q(x, \xi, t) = F_2^q(t) \quad \int dx \tilde{E}_q(x, \xi, t) = G_P^q(t),$$

- x_i and x_f are the momentum fractions of the quark before and after the momentum transfer
- $2\xi = x_f - x_i$
- GPDs can be probed in deeply virtual Compton scattering (DVCS)



Generalized Parton Distributions (GPDs)

- formal definition (unpol. quarks):

$$\int \frac{dx^-}{2\pi} e^{ix^- \bar{p}^+ x} \left\langle p' \left| \bar{q} \left(-\frac{x^-}{2} \right) \gamma^+ q \left(\frac{x^-}{2} \right) \right| p \right\rangle = H(x, \xi, \Delta^2) \bar{u}(p') \gamma^+ u(p) + E(x, \xi, \Delta^2) \bar{u}(p') \frac{i\sigma^{+\nu} \Delta_\nu}{2M} u(p)$$

- in the limit of vanishing t and ξ , the nucleon non-helicity-flip GPDs must reduce to the ordinary PDFs:

$$H_q(x, 0, 0) = q(x) \quad \tilde{H}_q(x, 0, 0) = \Delta q(x).$$

- DVCS amplitude

$$\mathcal{A}(\xi, t) \sim \int_{-1}^1 \frac{dx}{x - \xi + i\varepsilon} GPD(x, \xi, t)$$

Form Factors vs. GPDs

operator	forward matrix elem.	off-forward matrix elem.	position space
$\bar{q}\gamma^+q$	Q	$F(t)$	$\rho(\vec{r})$
$\int \frac{dx^-}{4\pi} e^{ixp^+x^-} \bar{q}\left(\frac{-x^-}{2}\right) \gamma^+ q\left(\frac{x^-}{2}\right)$	$q(x)$	$H(x, \xi, t)$?

Form Factors vs. GPDs

operator	forward matrix elem.	off-forward matrix elem.	position space
$\bar{q}\gamma^+q$	Q	$F(t)$	$\rho(\vec{r})$
$\int \frac{dx^-}{4\pi} e^{ixp^+x^-} \bar{q}\left(\frac{-x^-}{2}\right) \gamma^+ q\left(\frac{x^-}{2}\right)$	$q(x)$	$H(x, 0, t)$	$q(x, \mathbf{b}_\perp)$

$q(x, \mathbf{b}_\perp) =$ impact parameter dependent PDF

Impact parameter dependent PDFs

- define \perp localized state [D.Soper,PRD15, 1141 (1977)]

$$|p^+, \mathbf{R}_\perp = \mathbf{0}_\perp, \lambda\rangle \equiv \mathcal{N} \int d^2\mathbf{p}_\perp |p^+, \mathbf{p}_\perp, \lambda\rangle$$

Note: \perp boosts in IMF form Galilean subgroup \Rightarrow this state has

$$\mathbf{R}_\perp \equiv \frac{1}{P^+} \int dx^- d^2\mathbf{x}_\perp \mathbf{x}_\perp T^{++}(x) = \sum_i x_i \mathbf{r}_{i,\perp} = \mathbf{0}_\perp$$

(cf.: working in CM frame in nonrel. physics)

- define **impact parameter dependent PDF**

$$q(x, \mathbf{b}_\perp) \equiv \int \frac{dx^-}{4\pi} \langle p^+, \mathbf{R}_\perp = \mathbf{0}_\perp | \bar{q}(-\frac{x^-}{2}, \mathbf{b}_\perp) \gamma^+ q(\frac{x^-}{2}, \mathbf{b}_\perp) | p^+, \mathbf{R}_\perp = \mathbf{0}_\perp \rangle e^{ixp^+ x^-}$$





\hookrightarrow

$$\begin{aligned} q(x, \mathbf{b}_\perp) &= \int \frac{d^2\Delta_\perp}{(2\pi)^2} e^{i\Delta_\perp \cdot \mathbf{b}_\perp} H(x, 0, -\Delta_\perp^2), \\ \Delta q(x, \mathbf{b}_\perp) &= \int \frac{d^2\Delta_\perp}{(2\pi)^2} e^{i\Delta_\perp \cdot \mathbf{b}_\perp} \tilde{H}(x, 0, -\Delta_\perp^2), \end{aligned}$$

Impact parameter dependent PDFs

- No relativistic corrections (Galilean subgroup!)
- ↪ corollary: interpretation of 2d-FT of $F_1(Q^2)$ as charge density in transverse plane also free from relativistic corrections
- $q(x, \mathbf{b}_\perp)$ has probabilistic interpretation as number density ($\Delta q(x, \mathbf{b}_\perp)$ as difference of number densities)
- Reference point for IPDs is transverse center of (longitudinal) momentum $\mathbf{R}_\perp \equiv \sum_i x_i \mathbf{r}_{i,\perp}$
- ↪ for $x \rightarrow 1$, active quark ‘becomes’ COM, and $q(x, \mathbf{b}_\perp)$ must become very narrow (δ -function like)
- ↪ $H(x, 0, -\Delta_\perp^2)$ must become Δ_\perp indep. as $x \rightarrow 1$ (MB, 2000)
- ↪ consistent with lattice results for first few moments
- Note that this does not necessarily imply that ‘hadron size’ goes to zero as $x \rightarrow 1$, as separation \mathbf{r}_\perp between active quark and COM of spectators is related to impact parameter \mathbf{b}_\perp via $\mathbf{r}_\perp = \frac{1}{1-x} \mathbf{b}_\perp$.

$q(x, \mathbf{b}_\perp)$ for unpol. p



x = momentum fraction of the quark

$\vec{b} = \perp$ position of the quark

Transversely Deformed Distributions and $E(x, 0, -\Delta_{\perp}^2)$

M.B., Int.J.Mod.Phys.A18, 173 (2003)

- So far: only unpolarized (or long. pol.) nucleon! In general ($\xi = 0$):

$$\int \frac{dx^-}{4\pi} e^{ip^+ x^-} \langle P+\Delta, \uparrow | \bar{q}(0) \gamma^+ q(x^-) | P, \uparrow \rangle = H(x, 0, -\Delta_{\perp}^2)$$

$$\int \frac{dx^-}{4\pi} e^{ip^+ x^-} \langle P+\Delta, \uparrow | \bar{q}(0) \gamma^+ q(x^-) | P, \downarrow \rangle = -\frac{\Delta_x - i\Delta_y}{2M} E(x, 0, -\Delta_{\perp}^2).$$

- Consider nucleon polarized in x direction (in IMF)

$$|X\rangle \equiv |p^+, \mathbf{R}_{\perp} = \mathbf{0}_{\perp}, \uparrow\rangle + |p^+, \mathbf{R}_{\perp} = \mathbf{0}_{\perp}, \downarrow\rangle.$$


- ↪ unpolarized quark distribution for this state:

$$q(x, \mathbf{b}_{\perp}) = \mathcal{H}(x, \mathbf{b}_{\perp}) - \frac{1}{2M} \frac{\partial}{\partial b_y} \int \frac{d^2 \Delta_{\perp}}{(2\pi)^2} E(x, 0, -\Delta_{\perp}^2) e^{-i\mathbf{b}_{\perp} \cdot \Delta_{\perp}}$$

- Physics: $j^+ = j^0 + j^3$, and left-right asymmetry from j^3 !
[X.Ji, PRL 91, 062001 (2003)]

Intuitive connection with \vec{J}_q

- DIS probes quark momentum density in the infinite momentum frame (IMF). Quark density in IMF corresponds to $j^+ = j^0 + j^3$ component in rest frame (\vec{p}_{γ^*} in $-\hat{z}$ direction)
- $\hookrightarrow j^+$ larger than j^0 when quark current towards the γ^* ; suppressed when away from γ^*
- \hookrightarrow For quarks with positive orbital angular momentum in \hat{x} -direction, j^z is positive on the $+\hat{y}$ side, and negative on the $-\hat{y}$ side



- Details of \perp deformation described by $E_q(x, 0, -\Delta_\perp^2)$
- \hookrightarrow not surprising that $E_q(x, 0, -\Delta_\perp^2)$ enters Ji relation!

$$\langle J_q^i \rangle = S^i \int dx [H_q(x, 0, 0) + E_q(x, 0, 0)] x.$$

Transversely Deformed PDFs and $E(x, 0, -\Delta_{\perp}^2)$

- $q(x, \mathbf{b}_{\perp})$ in \perp polarized nucleon is deformed compared to longitudinally polarized nucleons !
- mean \perp deformation of flavor q (\perp flavor dipole moment)

$$d_y^q \equiv \int dx \int d^2 \mathbf{b}_{\perp} q_X(x, \mathbf{b}_{\perp}) b_y = \frac{1}{2M} \int dx E_q(x, 0, 0) = \frac{\kappa_q^p}{2M}$$

with $\kappa_{u/d}^p \equiv F_2^{u/d}(0) = \mathcal{O}(1 - 2) \Rightarrow d_y^q = \mathcal{O}(0.2 fm)$

- simple model: for simplicity, make ansatz where $E_q \propto H_q$

$$E_u(x, 0, -\Delta_{\perp}^2) = \frac{\kappa_u^p}{2} H_u(x, 0, -\Delta_{\perp}^2)$$
$$E_d(x, 0, -\Delta_{\perp}^2) = \kappa_d^p H_d(x, 0, -\Delta_{\perp}^2)$$






with $\kappa_u^p = 2\kappa_p + \kappa_n = 1.673$ $\kappa_d^p = 2\kappa_n + \kappa_p = -2.033$.

- Model too simple but illustrates that anticipated deformation is very significant since κ_u and κ_d known to be large!

p polarized in $+\hat{x}$ direction

$u(x, \mathbf{b}_\perp)$


$d(x, \mathbf{b}_\perp)$



lattice results (QCDSF)

GPD \longleftrightarrow SSA (Sivers)

- example: $\gamma p \rightarrow \pi X$



- u, d distributions in \perp polarized proton have left-right asymmetry in \perp position space (T-even!); sign “determined” by κ_u & κ_d
- attractive FSI deflects active quark towards the center of momentum
- \hookrightarrow FSI translates position space distortion (before the quark is knocked out) in $+\hat{y}$ -direction into momentum asymmetry that favors $-\hat{y}$ direction
- \hookrightarrow correlation between sign of κ_q^p and sign of SSA: $f_{1T}^{\perp q} \sim -\kappa_q^p$
- $f_{1T}^{\perp q} \sim -\kappa_q^p$ confirmed by HERMES data (also consistent with COMPASS deuteron data $f_{1T}^{\perp u} + f_{1T}^{\perp d} \approx 0$)

Quark-Gluon Correlations (Introduction)

- (longitudinally) polarized polarized DIS at leading twist \longrightarrow ‘polarized quark distribution’ $g_1^q(x) = q^\uparrow(x) + \bar{q}^\uparrow(x) - q_\downarrow(x) - \bar{q}_\downarrow(x)$
- $\frac{1}{Q^2}$ -corrections to X-section involve ‘higher-twist’ distribution functions, such as $g_2(x)$
- $g_2(x)$ involves quark-gluon correlations and does not have a parton interpretation as difference between number densities

Quark-Gluon Correlations (Introduction)

- (chirally even) higher-twist PDF $g_2(x) = g_T(x) - g_1(x)$

$$\int \frac{d\lambda}{2\pi} e^{i\lambda x} \langle PS | \bar{\psi}(0) \gamma^\mu \gamma_5 \psi(\lambda n) | Q^2 | PS \rangle$$
$$= 2 \left[g_1(x, Q^2) p^\mu (S \cdot n) + g_T(x, Q^2) S_\perp^\mu + M^2 g_3(x, Q^2) n^\mu (S \cdot n) \right]$$

- ‘usually’, contribution from g_2 to polarized DIS X-section kinematically suppressed by $\frac{1}{\nu}$ compared to contribution from g_1

$$\sigma_{TT} \propto g_1 - \frac{2Mx}{\nu} g_2$$

- for \perp polarized target, g_1 and g_2 contribute equally to σ_{LT}

$$\sigma_{LT} \propto g_T \equiv g_1 + g_2$$

- ↪ ‘clean’ separation between higher order corrections to leading twist (g_1) and higher twist effects (g_2)
- what can one learn from g_2 ?

Quark-Gluon Correlations (QCD analysis)

- (chirally even) higher-twist PDF $g_2(x) = g_T(x) - g_1(x)$

$$\int \frac{d\lambda}{2\pi} e^{i\lambda x} \langle PS | \bar{\psi}(0) \gamma^\mu \gamma_5 \psi(\lambda n) | Q^2 | PS \rangle$$

$$= 2 \left[g_1(x, Q^2) p^\mu (S \cdot n) + g_T(x, Q^2) S_\perp^\mu + M^2 g_3(x, Q^2) n^\mu (S \cdot n) \right]$$

- $g_2(x) = g_2^{WW}(x) + \bar{g}_2(x)$, with $g_2^{WW}(x) \equiv -g_1(x) + \int_x^1 \frac{dy}{y} g_1(y)$

- $\bar{g}_2(x)$ involves quark-gluon correlations, e.g.

$$\int dx x^2 \bar{g}_2(x) = \frac{1}{3} d_2 = \frac{1}{6MP^{+2}S^x} \langle P, S | \bar{q}(0) g G^{+y}(0) \gamma^+ q(0) | P, S \rangle$$

- $\sqrt{2}G^{+y} \equiv G^{0y} + G^{zy} = -E^y + B^x$

- matrix elements of $\bar{q}B^x\gamma^+q$ and $\bar{q}E^y\gamma^+q$ are sometimes called **color-electric and magnetic polarizabilities**

$$2M^2 \vec{S} \chi_E = \langle P, S | \vec{j}_a \times \vec{E}_a | P, S \rangle \quad \& \quad 2M^2 \vec{S} \chi_B = \langle P, S | j_a^0 \vec{B}_a | P, S \rangle$$

with $d_2 = \frac{1}{4} (\chi_E + 2\chi_M)$ — but **these names are misleading!**

Quark-Gluon Correlations (Interpretation)

- $\bar{g}_2(x)$ involves quark-gluon correlations, e.g.

$$\int dx x^2 \bar{g}_2(x) = \frac{1}{3} d_2 = \frac{1}{6MP^{+2}S^x} \langle P, S | \bar{q}(0) g G^{+y}(0) \gamma^+ q(0) | P, S \rangle$$

- QED: $\bar{q}(0) e F^{+y}(0) \gamma^+ q(0)$ correlator between quark density $\bar{q} \gamma^+ q$ and (\hat{y} -component of the) Lorentz-force

$$F^y = e \left[\vec{E} + \vec{v} \times \vec{B} \right]^y = e (E^y - B^x) = -e (F^{0y} + F^{zy}) = -e \sqrt{2} F^{+y}.$$

for charged particle moving with $\vec{v} = (0, 0, -1)$ in the $-\hat{z}$ direction

- ↪ matrix element of $\bar{q}(0) e F^{+y}(0) \gamma^+ q(0)$ yields γ^+ density (density relevant for DIS in Bj limit!) weighted with the Lorentz force that a charged particle with $\vec{v} = (0, 0, -1)$ would experience at that point
- ↪ d_2 a measure for the **color Lorentz force** acting on the struck quark in SIDIS in the instant after being hit by the virtual photon

$$\langle F^y(0) \rangle = -M^2 d_2 \quad (\text{rest frame; } S^x = 1)$$

Quark-Gluon Correlations (Interpretation)

- Interpretation of d_2 with the transverse FSI force in DIS also consistent with $\langle k_{\perp}^y \rangle \equiv \int_0^1 dx \int d^2 k_{\perp} k_{\perp}^2 f_{1T}^{\perp}(x, k_{\perp}^2)$ in SIDIS (Qiu, Sterman)

$$\langle k_{\perp}^y \rangle = -\frac{1}{2p^+} \left\langle P, S \left| \bar{q}(0) \int_0^{\infty} dx^- g G^{+y}(x^-) \gamma^+ q(0) \right| P, S \right\rangle$$

semi-classical interpretation: average k_{\perp} in SIDIS obtained by correlating the quark density with the transverse impulse acquired from (color) Lorentz force acting on struck quark along its trajectory to (light-cone) infinity

- matrix element defining d_2 same as the integrand (for $x^- = 0$) in the QS-integral:

- $\langle k_{\perp}^y \rangle = \int_0^{\infty} dt F^y(t)$ (use $dx^- = \sqrt{2}dt$)

↔ first integration point $\longrightarrow F^y(0)$

↔ (transverse) force at the begin of the trajectory, i.e. at the moment after absorbing the virtual photon

Quark-Gluon Correlations (Interpretation)

- x^2 -moment of twist-4 polarized PDF $g_3(x)$
$$\int dx x^2 g_3(x) \rightsquigarrow \langle P, S | \bar{q}(0) g \tilde{G}^{\mu\nu}(0) \gamma_\nu q(0) | P, S \rangle \sim f_2$$
 - ↪ different linear combination $f_2 = \chi_E - \chi_B$ of χ_E and χ_M
 - ↪ combine with data for $g_2 \Rightarrow$ disentangle electric and magnetic force
 - ↪ combining JLab(E99-117)/SLAC(E155x) data this yields
 - proton:
 $\chi_E = -0.082 \pm 0.016 \pm 0.071 \quad \chi_B = 0.056 \pm 0.008 \pm 0.036$
 - neutron:
 $\chi_E = 0.031 \pm 0.005 \pm 0.028 \quad \chi_B = 0.036 \pm 0.034 \pm 0.017$
- but future higher- Q^2 data for d_2 may still change these results ...

Quark-Gluon Correlations (Estimates)

- What should one expect (magnitude)?
 - if all spectators were to pull in the same direction, force should be on the order of the QCD string tension
 $\sigma \approx (0.45\text{GeV})^2 \approx 0.2\text{GeV}^2$
 - however, expect significant cancellation for FSI force, from spectators 'pulling' in different directions
 - ↪ expect FSI force to be suppressed compared to string tension by about one order of magnitude (more?)
 - ↪ $|d_2| = \frac{|\langle F^y(0) \rangle|}{M^2} \sim 0.02$
- What should one expect (sign)?
 - $\kappa_q^p \longrightarrow$ signs of deformation (u/d quarks in $\pm\hat{y}$ direction for proton polarized in $+\hat{x}$ direction \longrightarrow expect force in $\mp\hat{y}$)
 - ↪ d_2 positive/negative for u/d quarks in proton
 - d_2 negative/positive for u/d quarks in neutron
 - large N_C : $d_2^{u/p} = -d_2^{d/p}$
 - consistent with $f_{1T}^{\perp u} + f_{1T}^{\perp d} \approx 0$

Quark-Gluon Correlations (data/lattice)

- lattice (Göckeler et al.): $d_2^u \approx 0.010$ and $d_2^d \approx -0.0056$ (with large errors)

↪ using $M^2 \approx 5 \frac{\text{GeV}}{fm}$ this implies

$$\langle F_u^y(0) \rangle \approx -50 \frac{\text{MeV}}{fm} \qquad \langle F_d^y(0) \rangle \approx 28 \frac{\text{MeV}}{fm}$$

- signs consistent with impact parameter picture
- SLAC data (5GeV^2): $d_2^p = 0.007 \pm 0.004$, $d_2^n = 0.004 \pm 0.010$
- combined with SIDIS data for $\langle k^y \rangle$, should tell us about ‘effective range’ of FSI $R_{eff} \equiv \frac{\langle k^y \rangle}{F^y(0)}$
Anselmino et al.: $\langle k^y \rangle \sim \pm 100 \text{ MeV}$
- x^2 -moment of chirally odd twist-3 PDF $e(x) \longrightarrow$ **transverse force on transversely polarized quark in unpolarized target** (\leftrightarrow Boer-Mulders h_1^\perp)

Chirally Odd GPDs


$$\int \frac{dx^-}{2\pi} e^{ixp^+ x^-} \left\langle p' \left| \bar{q} \left(-\frac{x^-}{2} \right) \sigma^{+j} \gamma_5 q \left(\frac{x^-}{2} \right) \right| p \right\rangle = H_T \bar{u} \sigma^{+j} \gamma_5 u + \tilde{H}_T \bar{u} \frac{\varepsilon^{+j\alpha\beta} \Delta_\alpha P_\beta}{M^2} u + E_T \bar{u} \frac{\varepsilon^{+j\alpha\beta} \Delta_\alpha \gamma_\beta}{2M} u + \tilde{E}_T \bar{u} \frac{\varepsilon^{+j\alpha\beta} P_\alpha \gamma_\beta}{M} u$$

- See also M.Diehl+P.Hägler, EPJ C44, 87 (2005).
- Fourier trafo of $\bar{E}_T^q \equiv 2\tilde{H}_T^q + E_T^q$ for $\xi = 0$ describes distribution of transversity for unpolarized target in \perp plane

$$q^i(x, \mathbf{b}_\perp) = \frac{\varepsilon^{ij}}{2M} \frac{\partial}{\partial b_j} \int \frac{d^2 \Delta_\perp}{(2\pi)^2} e^{i\mathbf{b}_\perp \cdot \Delta_\perp} \bar{E}_T^q(x, 0, -\Delta_\perp^2)$$


- origin: correlation between quark spin (i.e. transversity) and angular momentum

Transversity Distribution in Unpolarized Target




IPDs on the lattice (QCDSF)

- lowest moment of distribution of unpol. quarks in \perp pol. proton (left) and of \perp pol. quarks in unpol. proton (right):



Transversity Distribution in Unpolarized Target (sign)

- Consider quark polarized out of the plane in ground state hadron
- ↪ expect counterclockwise **net current** \vec{j} associated with the magnetization density in this state



- virtual photon 'sees' enhancement of quarks (polarized out of plane) at the top, i.e.
- ↪ virtual photon 'sees' enhancement of quarks with polarization up (down) on the left (right) side of the hadron
- ↪ $\bar{E}_T > 0$

Transversity Distribution in Unpolarized Target (sign)

[M.B.+B.Hannafious, PLB 658, 1130 (2008)]

- matrix element for \bar{E}_T involves quark helicity flip
- ↪ requires interference between wave function components that differ by one unit of OAM (e.g. s-p interference)
- ↪ sign of \bar{E}_T depends on rel. sign between s & p components
- bag model: p-wave from lower component

$$\Psi_m = \begin{pmatrix} i f \chi_m \\ -g(\vec{\sigma} \cdot \hat{x}) \chi_m \end{pmatrix},$$

(relative sign from free Dirac equation $g = \frac{1}{E} \frac{d}{dr} f$)

- $\bar{E}_T \propto -f \cdot g$. Ground state: f peaked at $r = 0 \Rightarrow \bar{E}_T > 0$
- more general potential model: $\frac{1}{E} \rightarrow \frac{1}{E - V_0(r) + m + V_S(r)}$
- ↪ sign of \bar{E}_T same as in Bag model!

Boer-Mulders Function

- SIDIS: attractive FSI expected to convert position space asymmetry into momentum space asymmetry
 - ↪ e.g. quarks at negative b_x with spin in $+\hat{y}$ get deflected (due to FSI) into $+\hat{x}$ direction
 - ↪ (qualitative) connection between Boer-Mulders function $h_1^\perp(x, \mathbf{k}_\perp)$ and the chirally odd GPD \bar{E}_T that is similar to (qualitative) connection between Sivers function $f_{1T}^\perp(x, \mathbf{k}_\perp)$ and the GPD E .
- **Boer-Mulders**: distribution of \perp pol. quarks in unpol. proton

$$f_{q^\uparrow/p}(x, \mathbf{k}_\perp) = \frac{1}{2} \left[f_1^q(x, \mathbf{k}_\perp^2) - h_1^{\perp q}(x, \mathbf{k}_\perp^2) \frac{(\hat{\mathbf{P}} \times \mathbf{k}_\perp) \cdot S_q}{M} \right]$$

- $h_1^{\perp q}(x, \mathbf{k}_\perp^2)$ can be probed in DY (RHIC, J-PARC, GSI) and tagged SIDIS (JLab, eRHIC), using Collins-fragmentation


probing BM function in tagged SIDIS

- consider semi-inclusive pion production off unpolarized target
- spin-orbit correlations in target wave function provide correlation between (primordial) quark transversity and impact parameter
- ↪ (attractive) FSI provides correlation between quark spin and \perp quark momentum \Rightarrow BM function
- Collins effect: left-right asymmetry of π distribution in fragmentation of \perp polarized quark \Rightarrow 'tag' quark spin
- ↪ $\cos(2\phi)$ modulation of π distribution relative to lepton scattering plane
- ↪ $\cos(2\phi)$ asymmetry proportional to: Collins \times BM

probing BM function in tagged SIDIS


Primordial Quark Transversity Distribution

→ \perp quark pol.



\perp polarization and γ^* absorption


- QED: when the γ^* scatters off \perp polarized quark, the \perp polarization gets modified
 - gets reduced in size
 - gets tilted symmetrically w.r.t. normal of the scattering plane



probing BM function in tagged SIDIS

Primordial Quark Transversity Distribution


→ \perp quark pol.



probing BM function in tagged SIDIS

Quark Transversity Distribution after γ^* absorption

→ \perp quark pol.



quark transversity component in lepton scattering plane flips


probing BM function in tagged SIDIS

\perp momentum due to FSI

\rightarrow \perp quark pol.

\downarrow \mathbf{k}_{\perp}^q due to FSI

lepton scattering plane



on average, FSI deflects quarks towards the center

Collins effect

- When a \perp polarized struck quark fragments, the structure of jet is sensitive to polarization of quark
- distribution of hadrons relative to \perp polarization direction may be left-right asymmetric
- asymmetry parameterized by **Collins fragmentation function**
- Artru model:
 - struck quark forms pion with \bar{q} from $q\bar{q}$ pair with 3P_0 'vacuum' quantum numbers
 - ↪ pion 'inherits' OAM in direction of \perp spin of struck quark
 - ↪ produced pion preferentially moves to left when looking into direction of motion of fragmenting quark with spin up
- Artru model confirmed by HERMES experiment
- more precise determination of Collins function under way (BELLE)


probing BM function in tagged SIDIS

\perp momentum due to Collins

\mathbf{k}_\perp due to Collins
 \rightarrow \perp quark pol.

\downarrow \mathbf{k}_\perp^q due to FSI

lepton scattering plane



SSA of π in jet emanating from \perp pol. q

probing BM function in tagged SIDIS


net \perp momentum (FSI+Collins)

\downarrow \mathbf{k}_{\perp} due to Collins

\downarrow \mathbf{k}_{\perp}^q due to FSI

\downarrow net \mathbf{k}_{\perp}^q


lepton scattering plane



\hookrightarrow in this example, enhancement of pions with \perp momenta \perp to lepton plane

probing BM function in tagged SIDIS

net k_{\perp}^{π} (FSI + Collins)



↔ expect enhancement of pions with \perp momenta \perp to lepton plane

Quark-Gluon Correlations (chirally odd)

- \perp momentum for quark polarized in $+\hat{x}$ -direction (unpolarized target)

$$\langle k_{\perp}^y \rangle = \frac{g}{2p^+} \left\langle P, S \left| \bar{q}(0) \int_0^{\infty} dx^- G^{+y}(x^-) \sigma^{+y} q(0) \right| P, S \right\rangle$$

- compare: interaction-dependent twist-3 piece of $e(x)$

$$\int dx x^2 e^{int}(x) \equiv e_2 = \frac{g}{4MP^{+2}} \langle P, S | \bar{q}(0) G^{+y}(0) \sigma^{+y} q(0) | P, S \rangle$$

↪ $\langle F^y \rangle = M^2 e_2$

↪ (chromodynamic lensing) $e_2 < 0$

Summary

- GPDs \xleftrightarrow{FT} IPDs (impact parameter dependent PDFs)
- $E(x, 0, -\Delta_{\perp}^2) \longrightarrow \perp$ deformation of PDFs for \perp polarized target
- ↪ $\kappa^{q/p} \Rightarrow$ sign of deformation
- ↪ attractive FSI $\Rightarrow f_{1T}^{\perp u} < 0$ & $f_{1T}^{\perp d} > 0$
- Interpretation of $M^2 d_2 \equiv 3M^2 \int dx x^2 \bar{g}_2(x)$ as \perp force on active quark in DIS in the instant after being struck by the virtual photon

$$\langle F^y(0) \rangle = -M^2 d_2 \quad (\text{rest frame; } S^x = 1)$$

- In combination with measurements of f_2
 - color-electric/magnetic force $\frac{M^2}{4} \chi_E$ and $\frac{M^2}{2} \chi_M$
- $\kappa^{q/p} \Rightarrow \perp$ deformation $\Rightarrow d_2^{u/p} > 0$ & $d_2^{d/p} < 0$ (attractive FSI)
- combine measurement of d_2 with that of $f_{1T}^{\perp} \Rightarrow$ range of FSI
- x^2 -moment of chirally odd twist-3 PDF $e(x) \longrightarrow$ transverse force on transversely polarized quark in unpolarized target (\leftrightarrow Boer-Mulders h_1^{\perp})

Summary

- distribution of \perp polarized quarks in unpol. target described by chirally odd GPD $\bar{E}_T^q = 2\bar{H}_T^q + E_T^q$
- ↪ origin: correlation between orbital motion and spin of the quarks
- ↪ attractive FSI \Rightarrow measurement of h_1^\perp (DY, SIDIS) provides information on \bar{E}_T^q and hence on spin-orbit correlations

- expect:

$$h_1^{\perp,q} < 0 \qquad |h_1^{\perp,q}| > |f_{1T}^q|$$

- x^2 -moment of chirally odd twist-3 PDF $e(x) \longrightarrow$ **transverse force on transversely polarized quark in unpolarized target** (\longrightarrow Boer-Mulders)