



## INTERNATIONAL NUCLEAR PHYSICS CONFERENCE

ADELAIDE, AUSTRALIA  
ADELAIDE CONVENTION CENTRE  
11-16 September 2016

[www.inpc2016.com](http://www.inpc2016.com)




## Baryons Spectrum and Structure

Annalisa D'Angelo

University of Rome Tor Vergata & INFN Rome Tor Vergata  
Rome - Italy

### Outline:

- Why study spectroscopy
- Establishing  $N^*$  states
- Identifying the effective degrees of freedom
- Outlook & conclusions




# Why N\* ?

## Baryon Spectroscopy Reveals the Workings of QCD

*“Nucleons are the stuff of which our world is made.*

*As such they must be at the center of any discussion of why the world we actually experience has the character it does.”*

Nathan Isgur, NStar2000, Newport News,  
Virginia




Derek B. Leinweber – University of Adelaide



# Why N\* ? From the Hydrogen Spectrum to QCD



Niels Bohr (1922)



Spectral series of hydrogen



- Understanding the hydrogen atom's ground state requires understanding its excitation spectrum.


→ From Bohr model of the atom to QED.

- Understanding the proton's ground state requires understanding its excitation spectrum.


→ From the Constituent Quark model to QCD.

# Historical Markers

**1952:** First glimpse of the  $\Delta(1232)$  in  $\pi p$  scattering shows internal structure of the proton.




**1964:** Baryon resonances essential in establishing the quark model and the color degrees of freedom.




**1989:** Broad effort to address the missing baryon puzzle.





**2010:** First successful attempt to predict the nucleon spectrum in LQCD.



**2015:** Understanding of the baryon spectrum is needed to quantify the transition from QGP to the confined phase in the early universe.



# $N^*$ in the History of the Universe





Tetyana Galatyuk's talk

Dramatic events occur in the microsecond old Universe.

- The transition from the QGP to the baryon phase is dominated by excited baryons.  
A quantitative description requires more states than found to date => **missing baryons**.
- During the transition the quarks acquire **dynamical mass** and the **confinement of color** occurs.


# Constituent quark models and SU(6)xO(3)



- Current-quarks of perturbative QCD evolve into constituent quarks at low momentum.  
→ Connection between constituent and current quarks.
- QCD-inspired Constituent Quark models: states classified by isospin, parity and spin within each oscillator band. Many projected  $q^3$  states are still missing or uncertain.

# LQCD $N^*$ & $\Delta$ Spectra

- Exhibit the  $SU(6) \times O(3)$ -symmetry features
- Counting of levels consistent with non-rel. quark model
- Striking similarity with quark model
- No parity doubling




Problems are not solved!

Robert G. Edwards, Jozef J. Dudek, David G. Richards, Stephen J. Wallace  
Phys.Rev. D84 (2011) 074508




# What Do We Want to Learn ?

Understand the effective degrees of freedom underlying the  $N^*$  spectrum and the forces.




- A vigorous experimental program is worldwide underway with the aim to:
  - search for **undiscovered states** in meson **photoproduction** at CLAS, CBELSA, GRAAL, MAMI, LEPS
  - confirm or dismiss weaker candidates (\*, \*\*, \*\*\*)
  - characterize the  $N^*$  and  $\Delta$  spectrum systematics.
- Measure the strength of resonance excitations versus distance scale in meson **electro-production** at JLab, to reveal the **underlying degrees of freedom** in the  $Q^2$  evolution of the transition amplitudes.

# Establishing the $N^*$ and $\Delta$ Spectrum: $\pi N$ scattering




# Establishing the $N^*$ and $\Delta$ Spectrum: $\pi N$ Scattering


$d\sigma/d\Omega$



$P$



# Establishing the $N^*$ and $\Delta$ Spectrum: $\pi N$ Amplitudes





Isospin 1/2  
Imaginary T


Julia-Diaz, Lee, Matsuyama, Sato

# Establishing the $N^*$ and $\Delta$ Spectrum

Search all channels: not just  $\pi N$



## Photonuclear cross sections





# Establishing the $N^*$ and $\Delta$ Spectrum

Experimental requirements:

- Precision measurements of photo-induced processes in wide kinematics, e.g.  
 $\gamma p \rightarrow \pi N, \eta p, K\gamma, \dots$        $\gamma n \rightarrow \pi N, K^0\bar{K}^0, \dots$
- More complex reactions, e.g.  $\gamma p \rightarrow \omega p, p\phi, \pi\pi p, \eta\pi N, K^*\gamma, \dots$  may be sensitive to high mass states through direct transition to ground state or through cascade decays
- Polarization observables are essential

Engaging theoretical groups

Extract s-channel  
resonances




Hadronic  
production

&  
Electromagnetic  
production

# Polarization Observables: Complete Experiment

The holy grail of baryon resonance analysis



- Process described by **4** complex, parity conserving amplitudes
- **8** well-chosen measurements are needed to determine amplitude.
- Up to **16** observables measured directly
- **3** inferred from double polarization observables
- **13** inferred from triple polarization observables

David Ireland's talk

Beam ( $P^\gamma$ )	Target ( $P^T$ )	Recoil ( $P^R$ )			Target ( $P^T$ ) + Recoil ( $P^R$ )						
		$x'$	$y'$	$z'$	$x'$	$x'$	$x'$	$y'$	$y'$	$y'$	$z'$
unpolarized	$d\sigma_0$		$\hat{T}$			$\hat{P}$		$\hat{T}_{x'}$		$\hat{L}_{x'}$	$\hat{\Sigma}$
$P_L^\gamma \sin(2\phi_\gamma)$			$\hat{H}$	$\hat{G}$	$\hat{O}_{x'}$	$\hat{O}_{z'}$		$\hat{C}_{x'}$	$\hat{E}$	$\hat{F}$	$-\hat{C}_{x'}$
$P_L^\gamma \cos(2\phi_\gamma)$	$-\hat{\Sigma}$		$-\hat{P}$		$-\hat{T}$		$-\hat{L}_{z'}$	$\hat{T}_{z'}$	$-d\sigma_0$	$\hat{L}_{x'}$	$-\hat{T}_{x'}$
circular $P_c^\gamma$		$\hat{F}$		$-\hat{E}$	$\hat{C}_{x'}$	$\hat{C}_{z'}$		$-\hat{O}_{z'}$	$\hat{G}$	$-\hat{H}$	$\hat{O}_{x'}$


A. Sandorfi, S. Hoblit, H. Kamano, T.-S.H. Lee, J.Phys. 38 (2011) 053001

# Experimental set-up

Polarized Frozen-spin Targets & CEBAF Large Acceptance Spectrometer



or



+

Torus magnet

6 superconducting coils

start counter




Drift chambers

35,000 cells

Time-of-flight counters


plastic scintillators,  
684 photomultipliers



Electromagnetic calorimeters

Lead/scintillator, 1296 photomultipliers

Open CLAS detector



# CLAS N\* Experimental Program

	$\sigma$	$\Sigma$	T	P	E	F	G	H	$T_x$	$T_z$	$L_x$	$L_z$	$O_x$	$O_z$	$C_x$	$C_z$
$p\pi^0$	✓	✓	✓		✓	✓	✓	✓								
$n\pi^+$	✓	✓	✓			✓	✓	✓								
$p\eta$	✓	✓	✓			✓	✓	✓								
$p\eta'$	✓	✓	✓			✓	✓	✓								
$p\omega/\phi$	✓	✓	✓			✓	✓	✓								
$N\pi\pi$	✓	✓														
$K^+\Lambda$	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
$K^+\Sigma^0$	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
$K^0*\Sigma^+$	✓	✓									✓	✓				
$K^{+*}\Sigma^0$	✓	✓														
$p\pi^-$	✓	✓				✓	✓	✓								
$p\rho^-$	✓	✓				✓	✓	✓								
$K^-\Sigma^+$	✓	✓				✓	✓	✓								
$K^0\Lambda$	✓	✓		✓	✓	✓	✓	✓			✓	✓	✓	✓	✓	✓
$K^0\Sigma^0$	✓	✓		✓	✓	✓	✓	✓			✓	✓	✓	✓	✓	✓
$K^{0*}\Sigma^0$	✓	✓														

David Ireland's talk


Proton targets

Data taking completed May 18, 2012


✓-published, ✓-acquired

Neutron targets

# Establishing the $N^*$ and $\Delta$ Spectrum: $\gamma + p \rightarrow \pi^0 + p$




## $\Sigma$ Beam Asymmetry




Kamano  
Nakamura  
Lee &  
Sato, 2012

T single  
G E F double  
polarization  
observables  
also available


# Strangeness production: $\vec{\gamma} + \vec{p} \rightarrow K^+ + \bar{\Lambda} \rightarrow K^+ + p + \pi^-$




M. McCracken et al.(CLAS), Phys.RevC81,025201,2010



# Strangeness production: $\vec{\gamma} + \vec{p} \rightarrow K^+ + \bar{\Lambda} \rightarrow K^+ + p + \pi^-$



M. McCracken et al. (CLAS), Phys. Rev. C 81, 025201, 2010




A.V. Anisovich et al, EPJ A48, 15 (2012)

# The N(1900)3/2<sup>+</sup> State

- Bump first seen in SAPHIR  $K^+ \Lambda$  data but due to systematics in the data misinterpreted as  $J^P = 3/2^-$  (D-wave resonance).
- State was solidly established in Bn-Ga coupled-channel analysis making use of very precise  $K\Lambda$  polarized data, resulting in \*\*\* assignment in PDG2012. (P-wave resonance).
- State confirmed in an effective Langrangian resonance model analysis  $\gamma p \rightarrow K^+ + \Lambda$  (*O. V. Maxwell, PRC85, 034611, 2012*)
- State confirmed in a covariant isobar model single channel analysis  $\gamma p \rightarrow K^+ + \Lambda$  (*T. Mart & M. J. Kholili, PRC86, 022201, 2012*).
- First baryon resonance observed and multiply confirmed in electromagnetic production.



Candidate for \*\*\* state



# Updated Spectrum of Baryon Resonances

- From 2000 to 2010 no new Baryon resonances were considered by the PDG.
  - Used  $\pi N$ - scattering data and some  $\pi$ -photoproduction only.
- Mature multi-channel models now include many photoproduction data.
- E.g. Bonn-Gatchina PWA analysis, A. Anisovich et al. EPJ A 48, 15 (2012).


	Particle Data Group 2010	BnGa analyses	Particle Data Group 2012
N(1860)5/2 <sup>+</sup>		*	**
N(1875)3/2 <sup>-</sup>		***	***
N(1880)1/2 <sup>+</sup>		**	**
N(1895)1/2 <sup>-</sup>		**	**
N(1900)3/2 <sup>+</sup>	**	***	***
N(2060)5/2 <sup>-</sup>		***	**
N(2150)3/2 <sup>-</sup>		**	**
$\Delta(1940)3/2^-$	*	*	**

Naming scheme has changed:

$$L_{2I\ 2J}(E) \longrightarrow J^P(E)$$

- Results from photoproduction now add to the PDG tables and determine properties of baryon resonances


# Lower Mass $N^*/\Delta$ spectrum in 2015



Are there mass degenerate spin multiplets?  
Do these states fit into the SU(6) state symmetry? Lattice?



# Constituent Quark Models & QCD

SU(6)xO(3)




# Do New States Fit into Q<sup>3</sup> QM ?

SU(6)xO(3)




Quark Orbital Angular Momentum




# Do New States Fit into LQCD Projections ?

Ignoring the mass scale,  
new candidates fit the  $J^P$   
values predicted from  
LQCD.

The field would really  
benefit from more  
realistic Lattice masses  
for  $N^*$  states.




# Electroexcitation of $N^*/\Delta$ resonances



# Total cross section at $W < 2.1$ GeV




Data: K. Park et al. PRC 77 (2008) 015208; K. Park et al. PRC 91 (2015) 045203



Analysis with UIM & fixed- $t$  DR; Recent review: I. Aznauryan, V. Burkert, Prog. Part. Nucl. Phys. 67 (2012) 1.

# Electroexcitation kinematics



$$\frac{d^4\sigma}{dQ^2 dW d\Omega_K} = \Gamma(Q^2, W) \times \frac{d\sigma}{d\Omega_K}(Q^2, W, \Theta_K, \epsilon, \phi)$$

Virtual  
photon  
flux

Electroproduction  
cross section

$$\frac{d\sigma}{d\Omega_K} = \sigma_T + \epsilon_L \sigma_L + \epsilon \sigma_{TT} \cos(2\phi) + \sqrt{2\epsilon_L(\epsilon+1)} \sigma_{LT} \cos(\phi) + h\sqrt{2\epsilon_L(1-\epsilon)} \sigma_{LT'}$$

Transverse

Transverse-tra interference

e

Helicity structure


$\sigma_u$   
"Unseparated"

Longitudinal (sensitive to  $J=0^\pm$  exchange in t-channel: mesons, diquarks)

Transverse-longitudinal interference

Measured  $\sigma$  are decomposed using UIM or fixed-t DR to extract  $N^*$  &  $\Delta$  helicity amplitudes.

# Electrocouplings of the ‘Roper’ in 2002



$N(1440)1/2^+$



In 2002 Roper amplitude  $A_{1/2}$  measurements were more consistent with hybrid state  
but data were limited with large uncertainties.

# Electrocouplings of the ‘Roper’ in 2012

I. Aznauryan et al. (CLAS), PRC 80, 055203 (2009)

V. Mokeev et al. (CLAS), PRC 86, 035203 (2012)

N(1440) $1/2^+$



L. Tiator et al., Chin Phys C33, (2009) 1069 (MAID fit)

I. Aznauryan et al. PRC 76, (2007) 025212

Z.P. Li et al. PRD 46, (1992) 70

I. T. Obukhovsky et al. PRD 84, (2011) 014004

# Electrocouplings of the ‘Roper’ in 2016

N $\pi$  loops to model MB cloud: **running quark mass** in LF RQM.  
I. G. Aznauryan, V.D. Burkert PR C 85 (2012) 055202.


No loops to model MB cloud in LF RQM: **frozen constituent quark mass** in LF RQM .

I. T. Obukhovsky et al. PRD 89, (2014) 0140032.

Quark-core contributions from DSE/QCD  
J. Segovia et al. PRL 115 (2015) 171801.

**Meson Baryon cloud** inferred from CLAS data as the difference between data and the quark-core evaluation in DSE/QCD.



V. Mokeev et al., PR C 93 (2016) 025206.



The structure of the Roper is driven by the interplay of the core of three dressed quarks in the 1<sup>st</sup> radial excitation and the external meson-baryon cloud.

# MB Contribution to electro-excitation of N(1535)1/2-

Is it a 3-quark state or a hadronic molecule?





- Meson-baryon cloud may account for discrepancies at low  $Q^2$ .

# MB Contribution to electro-excitation of N(1675)5/2<sup>-</sup>

Quark components to the helicity amplitudes of the N(1675) 5/2<sup>-</sup> are strongly suppressed for proton target.

Single Quark Transition:  
 $A_{1/2}^p = A_{3/2}^p = 0$



- Measures the meson-baryon contribution to the  $\gamma^* p$  N(1675)5/2<sup>-</sup> directly.
- Can be verified on  $\gamma^* n$  N(1675)5/2<sup>-</sup> which is not suppressed

— E. Santopinto and M. M. Giannini, PRC 86, 065202 (2012)  
- - - B. Juliá-Díaz, T.-S.H. Lee, et al., PRC 77, 045205 (2008)

# Hybrid Baryons: Baryons with Explicit Gluonic Degrees of Freedom

Hybrid hadrons with dominant gluonic contributions are predicted to exist by QCD.

Experimentally:

- **Hybrid mesons**  $|q\bar{q}g\rangle$  states may have exotic quantum numbers  $J^{PC}$  not available to pure  $|q\bar{q}\rangle$  states      GlueX, MesonEx, COMPASS, PANDA ....
- **Hybrid baryons**  $|qq\bar{q}g\rangle$  have the same quantum numbers  $J^P$  as  $|qqq\rangle$  electroproduction with CLAS12 (Hall B).


Hugh Montgomery's talk

Theoretical predictions:

- ✧ MIT bag model - T. Barnes and F. Close, Phys. Lett. 123B, 89 (1983).
- ✧ QCD Sum Rule - L. Kisslinger and Z. Li, Phys. Rev. D 51, R5986 (1995).
- ✧ Flux Tube model - S. Capstick and P. R. Page, Phys. Rev. C 66, 065204 (2002).
- ✧ LQCD - J.J. Dudek and R.G. Edwards, PRD85, 054016 (2012).

# Hybrid Baryons in LQCD

J.J. Dudek and R.G. Edwards, PRD85, 054016 (2012)




Hybrid states have same  $J^P$  values as qqq baryons. How to identify them?

- Overpopulation of  $N \frac{1}{2}^+$  and  $N \frac{3}{2}^+$  states compared to QM projections.
- $A_{1/2}$  ( $A_{3/2}$ ) and  $S_{1/2}$  show different  $Q^2$  evolution.

# Separating $q^3g$ from $q^3$ states ?

CLAS results on electrocouplings clarified nature of the Roper.

Will CLAS12 data be able to identify gluonic contributions ?



For hybrid “Roper”,  $A_{1/2}(Q^2)$  drops off faster with  $Q^2$  and  $S_{1/2}(Q^2) \sim 0$ .

# Baryon Spectroscopy Status Today


- Major progress made in the last ~ 5 years in the search for  $N^*$  and  $\Delta$  states.  
All states can be accommodated in CQM and LQCD schemes.
  - Naïve (non-dynamical) di-quark models are ruled out.
- Knowledge of  $Q^2$ -dependence of electrocouplings is absolutely necessary to understand the nature ( the internal structure) of the excited states.
  - Roper IS the first radial excitation of the  $q^3$  core, obscured at large distances by meson-cloud effects.
- Leading electrocoupling amplitudes of prominent low-mass states (e.g.  $N(1535)1/2^-$ ) is well modeled by DSE/QCD, LC SR and LF RQM for  $Q^2 > 2$  GeV.
- Search for hybrid baryons with explicit gluonic degrees of freedom would be possible investigating the low  $Q^2$  evolution of high-mass resonance (2-3 GeV) electrocouplings:
  - Looking for suppressed  $A^{1/2}$ ,  $A^{3/2}$ ,  $S^{1/2}$  at low  $Q^2$ .

# Thank you !

# BACKUP SLIDES

# Missing Baryons in QCD Phase Transition

From Hot QCD:  
Fluctuation Ratio  
of Baryon Number  
to Strangeness at  
hadron freeze-out



A. Bazavov et al.,  
Phys.Rev.Lett. 113  
(2014) 7, 072001

Transition shifted  
by about 8 MeV to  
lower temperature  
(later times) due to  
missing excited  
strange baryons

→ The number of known excited strange baryon states (PDG) is insufficient to account for the QCD phase cross-over from the QGP phase to the baryon phase.

- Evidence for experimentally-missing strange baryons
- Evidence observed also for missing charm and light quark baryons
- Motivates an excited baryon program of all quark flavors.

The RHIC operation plan for 2016 includes an energy scan to map out this behavior.